

บรรณานุกรม

- บริษัท กิบไทย จำกัด. (2015). Cell Proliferation & Metabolic Activity. สืบค้นจาก http://www.gibthai.com/service/note_detail/41
- รัชฎาพร อุ่นศิวิไลย์. (2554). รายงานวิจัยเรื่อง ฤทธิ์ทางชีวภาพและคุณสมบัติเชิงหน้าที่ของสารสกัดย่านาง เครื่อหมาน้อย และรังจีด. มหาวิทยาลัยเทคโนโลยีสุรนารี นครราชสีมา
- Adler-Nissen J. (1979). Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzenesulfonic acid. *Journal of Agricultural and Food Chemistry*, 27, 1256-1262.
- Al-Dhabi, N.A., Ponnurugan, K., & Jeganathan, P.M. (2017). Development and validation of ultrasound-assisted solid-liquid extraction of phenolic compounds from waste spent coffee grounds. *Ultrasonics Sonochemistry*, 34, 206-213.
- Alam, M.N., Bristi, N.J., & Rafiquzzaman, M. (2013). Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*, 21, 143-152.
- Andrews, A. L., Atkinson, D., Evans, M. T. A., Finer, E. G., Green, J. P., Phillips, M. C., & Robertson, R. N. (1979). The conformation and aggregation of bovine β -casein. A. I. Molecular aspects of thermal aggregation. *Biopolymers*, 18, 1105-1121
- Ballesteros, L.F., Cerqueira, M.A., Teixeira, J.A., & Mussatto, S.I. (2015). Characterization of polysaccharides extracted from spent coffee grounds by alkali pretreatment. *Carbohydrate Polymers*, 127, 347-354.
- Ballesteros, L.F., Teixeira, J.A., & Mussatto, S.I. (2014). Chemical, functional, and structural properties of spent coffee grounds and coffee silverskin. *Food and Bioprocess Technology*, 7, 3493-3503.
- Ballesteros, L.F., Teixeira, J.A., & Mussatto, S.I. (2017). Extraction of polysaccharides by autohydrolysis of spent coffee grounds and evaluation of their antioxidant activity. *Carbohydrate Polymers*, 157, 258-266.
- Bravo, J., Arbillage, L., Peña, M.P., & Cid, C. (2013). Antioxidant and genoprotective effects of spent coffee extracts in human cells. *Food and Chemical Toxicology*, 60, 397-403.
- Bravo, J., Juániz, I., Monente, C., Caemmerer, B., Kroh, L.W., Peña, M.P., & Cid, C. (2012). Evaluation of spent coffee obtained from the most common coffeemakers as

- source of hydrophilic bioactive compounds. *Journal of Agricultural and Food Chemistry*, 60, 12565-12573.
- Bravo, J., Monente, C., Juániz, I., Peña, M.P., & Cid, C. (2013). Influence of extraction process on antioxidant capacity of spent coffee. *Food Research International*, 50, 610-616.
- Cämmerer, B., & Kroh, L.W. (2006). Antioxidant activity of coffee brews. *European Food Research and Technology*, 223, 469-474
- Castillo, M, D., Ames, J.M., & Gordon, A, H. (2002). Effect of roasting on the antioxidant activity of coffee brews. *Journal of Agricultural and Food Chemistry*, 50, 3698-3703.
- Cha, K.H., Song, D.-G., Kim, S.-M., & Pan, C.-H. (2012). Inhibition of gastrointestinal lipolysis by green tea, coffee, and gomchui (*Ligularia fischeri*) tea polyphenols during simulated digestion. *Journal of Agricultural and Food Chemistry*, 60, 7152-7157.
- Chen, M.-F., Zhang, Y.Y., He, M.D., Li, C.Y., Zhou, C.X., Hong, P.Z., & Qian, Z.-J. (2019). Antioxidant Peptide Purified from Enzymatic Hydrolysates of Isochrysis Zhanjiangensis and Its Protective Effect against Ethanol Induced Oxidative Stress of HepG2 Cells. *Biotechnology and Bioprocess Engineering*, 24, 308-317.
- Chi, C., Li, X., Miao, S., Zhang, Y., & Chen, L.(2019). Dry heating and annealing treatment synergistically modulate starch structure and digestibility. *International Journal of Biological Macromolecules*, 137, 554-561.
- Chatthongpisut, R., Schwartz, S.J., and Yongsawatdigul, J. (2015). Antioxidant activities and antiproliferation activity of Thai purple rice cooked by various methods on human colon cancer cells. *Food Chemistry*, 188, 99-105.
- Charupin, P., Ames, J.M, & Castillo, M, D. (2002). Antioxidant activity of coffee model systems. *Journal of Agricultural and Food Chemistry*, 50, 3751-3756.
- Choi, B., & Koh, E. (2017). Spent coffee as a rich source of antioxidative compounds. *Food Science and Biotechnology*, 26, 921-927.
- Compos-Vega, R., Loarca-Piña, G., Vergara-Castañeda, H.A., and Oomah, B.D. (2015). Spent coffee grounds: A review on current research and future prospects. *Trends in Food Sciences & Technology*, 45, 24-36.

- Compos-Vega, R., Vázquez-Sánchez, K., López-Barrera, D., Loarca-Piña, G., Mendoza-Díaz, S., & Oomah, B.D. (2015). Simulated gastrointestinal digestion and in vitro colonic fermentation of spent coffee (*Coffea arabica* L.): bioaccessibility and intestinal permeability. *Food Research International*, 77, 156-161.
- Cruz, R., Caroso, M.M., Fernandes, L., Oliveira, M., Mendes, E., Baptista, P., Morais, S., & Casal, S. (2012). Espresso coffee residues: a valuable source of unextracted compounds. *Journal of Agricultural and Food Chemistry*, 60, 7777-7784.
- Daglia, M., Papetti, A., Gregotti, C., Bertè, F., & Gazzani, G. (2000). In vitro antioxidant and ex vivo protective activities of green and roasted coffee. *Journal of Agricultural and Food Chemistry*, 48, 1449-1454.
- Dias, M.I., Morales, P., Barreira, J.C.M., Oliveira, M.B.P.P., Sanchez-Mata, M.C., & Ferreira, I.C.F.R. (2016). Minerals and vitamin B9 in dried plants vs. infusions: Assessing absorption dynamics of minerals by membrane dialysis tandem in vitro digestion. *Food Bioscience*, 13, 9-14.
- Decker E.A., & Welch, B. (1990). Role of ferritin as a lipid oxidation catalyst in muscle food. *Journal of Agricultural and Food Chemistry*, 38, 674-677.
- Delgado-Andrade, C., Rufián-Henares, J.A., & Morales, F.J. (2005). Assessing the antioxidant activity of melanoidins from coffee brews by different antioxidant methods. *Journal of Agricultural and Food Chemistry*, 53, 7832-7836.
- Delgado-Andrade, C., & Morales, F.J. (2005). Unraveling the contribution of melanoidins to the antioxidant activity of coffee brews. *Journal of Agricultural and Food Chemistry*, 53, 1403-1407.
- Duarté, S.M.S., de Abreu, C.M.P., Menezes, H.C. dos Santos, M.H., & Gouvêa, C.M.C.P. (2005). Effect of processing and roasting on the antioxidant activity of coffee brews. *Ciência e Tecnologia de Alimentos*, 25, 387-393.
- Gómez-Ruiz, J.A., Leake, D.S., & Ames, J.M. (2007). In vitro antioxidant activity of coffee compounds and their metabolites. *Journal of Agricultural and Food Chemistry*, 55, 6962-6969.
- International Coffee Organization (2020). Trade statistics tables. Available at <http://www.ico.org/prices/po-production.pdf>

- Iriondo-DeHond, A., Ramírez, B., Escobar, F.V., & and del Castillo1, M.D. (2019). Antioxidant properties of high molecular weight compounds from coffee roasting and brewing byproducts. *Bioactive Compounds in Health and Disease*, 2, 48-63.
- Jin, Z., Bai, F., Chen, Y. & Bai, B. (2016). Interactions between protein, lipid and starch in foxtail millet flour affect the in vitro digestion of starch. *CyTA - Journal of Food*, 17, 640-647.
- Jully, K.M.M., Tota, C.S., & Were, L. (2016). Antioxidant effect of spent, ground, and lyophilized brew from roasted coffee in frozen cooked pork patties. *LWT-Food Science and Technology*, 66, 244-251.
- Krishnaiah, D., Sarbatly, R., & Nithyanandam, R. (2011). A review of antioxidant potential of medicinal plant species. *Food and Bioproducts Processing*, 89, 217-233.
- López-Barrera, D.M., Vázquez-Sánchez, Loarca-Piñn, G, F., & Campos-Vega, R. (2016). Spent coffee grounds, an innovative source of colonic fermentable compounds, inhibit inflammatory mediators *in vitro*. *Food Chemistry*, 212, 282-290.
- Ludwing, I., Sanchez, L., Caemmerer, B., Kroh, L.W., Peña, M.P., & Cid, C (2012). Extraction of coffee antioxidants: impact of brewing time and method. *Food Research International*, 48, 57-64.
- Martinez-Saez, N., García, A.T., Pérez, I.D., Rebollo-Hernanz, M., Mesías, M., Morales, F.J., Martín-Cabrejas, M.A., & del Castillo, M.D. (2017). Use of spent coffee grounds as food ingredient in bakery products. *Food Chemistry*, 216, 114–122.
- Mineku et al. (2014) A standardised static in vitro digestion method suitable for food – an international consensus. *Food and Function*, 5, 1113-1124
- Monnente, C., Ludwig, I.A., Irigoyen, A., Peña, M.P., & Cid, C. (2015). Assessment of total (free and bound) phenolic compounds in spent coffee extracts. *Journal of Agricultural and Food Chemistry*, 63, 4327-4334.
- Moon, J.;K., Yoo, H.S., & Shibamoto, T. (2009). Role of roasting conditions in the level of chlorogenic acid content in coffee beans: correlation with coffee acidity. *Journal of Agricultural and Food Chemistry*, 57, 5365-5369.

- Musatto, S., Ballesteros, L.F., Martins, S., and Teixeria, J.A. (2011). Extraction of antioxidant phenolic compounds from spent coffee grounds. *Separation and Purification Technology*, 83, 173-179.
- Nicoli, M.C., Anese, M., Manzocco, L., & Lerici, C.R. (1997). Antioxidant properties of coffee brews in relation to the roasting degree. *LWT - Food Science and Technology*, 30, 292-297.
- Nimse, S.B., & Pal, D. (2015). Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Advances*, 5, 27986
- Panzella, L., Cerruti, P., Ambrogi, V., Agustin-Salazar, S., D'Errico, G., Cosimo Carfagna, C., Goya, L., Ramos, S., Martín, M.A., Napolitano, A., & d'Ischia, M. (2016). A superior all-natural antioxidant biomaterial from spent coffee grounds for polymer stabilization, cell protection, and food lipid preservation. *ACS Sustainable Chemistry & Engineering*, 4, 1169-1179.
- Pérez-Cruz, K., Moncada-Basualto, M., Morales-Valenzuela, J., Barriga-González, G., Navarrete-Encina, P., Núñez-Vergara, L., Squella, J.A., & Olea-Azar, C. (2018). Synthesis and antioxidant study of new polyphenolic hybrid-coumarins. *Arabian Journal of Chemistry*, 11, 525-537.
- Quirós-Sauceda, A. E., Palafox-Carlos, H., Sáyago-Ayerdi, S. G., Ayala-Zavala, J. F., Bello-Perez, L. A., Álvarez-Parrilla, E. , de la Rosa, L. A., González-Córdovaa, A. F., & González-Aguilar, G. A. . (2014). Dietary fiber and phenolic compounds as functional ingredients: interaction and possible effect after ingestion. *Food and Function*, 5, 1063-1072
- Ramalakshmi, K., Rao, L.J.M., Takano-Ishikawa, Y., & Goto, M. (2009). Bioactivities of low-grade green coffee and spent coffee in different in vitro model systems. *Food Chemistry*, 115, 79-85.
- Ranic, M., Nikolic, M., Pavlovic, M., Buntic, A., Siler-Marinkovic, S., & Dimitrijevic-Brankovic, S. (2014). Optimization of microwave-assisted extraction of natural antioxidants from spent espresso coffee grounds by response surface methodology. *Journal of Cleaner Production*, 80, 69-79.

- Re, R., Pellegrini, N., Proteggente, A., Pannalla, A., Yang, M., and Rice-Evans, C. (1995). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 53, 1231-1237.
- Richelle, M., Tavazzi, I., & Offord, E. (2001). Comparison of the antioxidant activity of commonly consumed polyphenolic beverage (coffee, cocoa, and tea) prepared per cup serving. *Journal of Agricultural and Food Chemistry*, 49, 3438-3442.
- Rufián-Henares, J.A. & Morales, F.J. (2007). Effect of in vitro enzymatic digestion on antioxidant activity of coffee melanoidins and fractions. *Journal of Agricultural and Food Chemistry*, 55, 10016-10021.
- Sam, R. (2014). Hemodialysis: Diffusion and ultrafiltration. *Austin Journal of Nephrology and Hypertension*, 1, 1010.
- Sánchez-González, I., Jiménez-Escríg, A., & Saura-Calixto, F. (2005). In vitro antioxidant activity of coffees brewed using different procedures (Italian, espresso and filter). *Food Chemistry*, 90, 133-139.
- Sangsawad, P., Kiatsongchai, R., Chitsomboon, B., & Yongsawatdigul, J. (2016). Chemical and cellular antioxidant activities of chicken breast muscle subjected to various thermal treatments followed by simulated gastrointestinal digestion. *Journal of Food Science*, 81, C2431-C2438
- Shao, Y., & Lin, A. H.-M. (2018). Improvement in the quantification of reducing sugars by miniaturizing the Somogyi-Nelson assay using a microtiter plate. *Food chemistry*, 240, 898–903
- The Refresher Co., Ltd. (2009). ກາແພ.ສືບຄໍນຈາກ http://www.refresherthai.com/article/species_of_coffee.php
- Ustunol, Z. (2015). Amino acids, peptides, and proteins. In Z. Ustunol (Ed.), *Applied Food Chemistry* (pp. 11-21). Singapore: Wiley Blankwell.
- Vignoli, J.A., Viegas, M.C., Bassoli, D.G., & Benassi, M. T. (2014). Roasting process affects differently the bioactive compounds and the antioxidant activity of arabica and robusta coffees. *Food Research International*, 61, 279-258.
- Wang, L., Li, X., Li, Y., Liu, W., Jia, X., Qiao, X., Qu, C., Cheng, X., & Wang, S. (2018). Antioxidant and angiotensin I-converting enzyme inhibitory activities of Xuanwei

- ham before and after cooking and in vitro simulated gastrointestinal digestion. *Royal Society Open Science*, 5, 180276.
- Xie, N., Wang, C., Ao, J., & Li, B. (2013). Non-gastrointestinal-hydrolysis enhances bioavailability and antioxidant efficacy of casein as compared with its in vitro gastrointestinal digest. *Food Research International*, 51, 114–122.
- Xu, H., Wang, W.W., Liu, X., Yuan, F., & Gao, Y. (2015). Antioxidative phenolics obtained from spent coffee grounds (*Coffea arabica* L.) by subcritical water extraction. *Industrial Crops and Products*, 76, 946-954.
- Yilmaz, Y., & Toledo, R. (2005), Antioxidant activity of water-soluble Maillard reaction products. *Food Chemistry*, 93, 273-278.
- Zhang, Z., Zhang, R., Xiao, H., Bhattacharya, K. Bitounis, D., Demokritou, P., and MaClements, D.J. (2019). Development of a standardized food model for studying the impact of food matrix effects on the gastrointestinal fate and toxicity of ingested nanomaterials. *NanoImpact*. 13, 13-25.
- Zhou, N., Zhu, W., Yang, F., & Zhou, K. (2016). In vitro gastrointestinal digestion model to monitor the antioxidant properties and bioavailability of phenolic antioxidants from elderberry. *Reactive Oxygen Species*, 2, 421-431.